

Product datasheet for AP06757PU-S

PERK (EIF2AK3) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunofluorescence: 1/50-1/200.

Reactivity: Human Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 56-102 of Human PERK.

Specificity: This antibody detects endogenous levels of PERK protein.

(region surrounding Arg87)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Preservative: 15 mM Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 125 kDa

Gene Name: eukaryotic translation initiation factor 2 alpha kinase 3

Database Link: Entrez Gene 9451 Human

Q9NZ<u>J5</u>



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:


An interferon-inducible, RNA-dependent protein serine /threonine kinase (PKR) has been described. PKR in earlier literature is variously known as DAI, dsJ, PI kinase, p65, p67 or TIK for the mouse kinase; and p68 or p69 for the human kinase. The PKR kinase substrate is the alpha subunit of protein synthesis initiation factor eIF-2. Phosphorylation of eIF-2alpha on serine-51 results in inhibition of translation. The serine /threonine kinase catalytic domains map to the carboxy terminal half of the protein while the RNA-binding domains are located in the amino terminal region. PERK is a type I transmembrane protein located in the endoplasmic reticulum (ER) that contains a kinase domain similar to the kinase domain of PKR. PERK is activated in response to ER stress and phosphorylates eIF-2alpha, thus inhibiting the translation of mRNA.

Synonyms: PEK, PERK, HsPEK

Product images:



Western blot (WB) analysis of PERK antibody in extracts from HeLa cells.



Immunohistochemistry analysis of PERK antibody on paraffin-embedded sections