

Product datasheet for **AP06758PU-N**

BRAF35 (HMG20B) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. Immunofluorescence: 1/50-1/200.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to the N-terminus of Human BRAF35.
Specificity:	This antibody detects endogenous levels of BRAF35 protein. (region surrounding Thr29)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 36 kDa
Gene Name:	high mobility group 20B
Database Link:	Entrez Gene 10362 Human Q9P0W2




[View online »](#)

Background:

The breast cancer susceptibility gene (BRCA1) localizes to chromosome 17q. Mutations within this gene account for approximately 45% of families with high incidence of breast cancer and at least 80% of families with increased incidence of both early-onset breast cancer and ovarian cancer. A second breast cancer susceptibility gene, BRCA2, located on chromosome 13q12-13, also confers a high incidence of breast cancer, but unlike BRCA1, BRCA2 does not confer a substantially elevated risk of ovarian cancer. The BRCA2-Associated Factor 35 (BRAF35) protein forms a 20 kDa complex with BRCA2, which associates with condensed chromatin during histone H3 phosphorylation. BRAF35 expression levels are highest in proliferating tissues and parallel BRCA2 expression patterns. The structure of BRAF35 includes a kinesinlike coiled coil domain and a nonspecific DNA binding HMG domain. The chromatin localization of BRAF35 and antibody microinjection studies indicate a role for the BRAF35/BRCA2 complex in cell cycle regulation.

Synonyms:

HMG20B, BRAF35, HMGX2, HMGXB2, SMARCE1R

Product images:

Western blot (WB) analysis of BRAF35 antibody at 1/500 dilution. Lane 1: A549 whole cell lysate; Lane 2: Mouse liver tissue lysate; Lane 3: Rat liver tissue lysate.