

Product datasheet for AP06807PU-N

XPF (ERCC4) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

ELISA, WB **Applications:**

Recommended Dilution: Western blot: 1/500-1/1000.

Reactivity: Human, Mouse

Rabbit Host:

Clonality: Polyclonal

Synthetic peptide, corresponding to amino acids 796-840 of Human ERCC4. Immunogen:

Specificity: This antibody detects endogenous levels of ERCC4/XPF protein.

Formulation: Phosphate buffered saline (PBS), pH~7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% by SDS-PAGE)

Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~117 kDa

Gene Name: ERCC excision repair 4, endonuclease catalytic subunit

Database Link: Entrez Gene 2072 Human

Q92889



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Xeroderma Pigmentosum (XP) is an autosomal recessive disorder characterized by a genetic predisposition to sunlight-induced skin cancer, and it is commonly due to deficiencies in DNA repair enzymes. The most frequent mutations are found in the XP genes from group A through G and group V, which encode for nucleotide excision repair proteins. XPF, which is also designated ERCC4 or ERCC11, is a 115 kDa protein that associates directly with the excision repair cross-complementing 1 (ERCC1) factor. ERCC-1, a functional homolog of Rad10 in S. cerevisiae, is a component of a structure-specific endonuclease that is responsible for 5' incisions during DNA repair. The ERCC1-XPF endonuclease preferentially cleaves one strand of DNA between duplex and singlestranded regions near borders of the stem-loop structure and, thereby, contributes to the initial steps of the nucleotide excision repair process.

Synonyms: ERCC-4, ERCC11, XPF

Product images:



Western blot analysis of ERCC4 Antibody in extracts from 293 cells.