

Product datasheet for **AP08015PU-N**

HSF1 pSer307 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, WB
Recommended Dilution:	Western blot (1/500-1/1000). Immunofluorescence (1/100-1/200).
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	The antiserum was produced against synthesized phosphopeptide derived from human HSF1 around the phosphorylation site of serine 307 (P-Q-SP-P-R).
Specificity:	This antibody HSF1 (pSer307) detects endogenous levels of HSF1 only when phosphorylated at Serine 307.
Formulation:	PBS (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol. State: Aff - Purified State: Liquid purified Ig fraction.
Concentration:	lot specific
Purification:	Immunoaffinity Chromatography: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.
Conjugation:	Unconjugated
Storage:	Store the antibody (in aliquots) at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	heat shock transcription factor 1
Database Link:	Entrez Gene 3297 Human Q00613



[View online »](#)

Background:

All organisms respond to elevated temperatures and a variety of environmental stresses by rapid synthesis of heat shock RNAs and proteins. The regulation of heat shock gene transcription is mediated by the transcriptional activator, heat shock factor (HSF), which binds to heat shock response elements (HSEs). These HSEs are found as three repeats of a 5-nucleotide {nGAAn} module, arranged in alternating orientation and present upstream of all heat shock genes. The HSEs are highly conserved among species yet HSF purified from yeast, *Drosophila* and human have different molecular weights and the proteins do not show significant immunological cross reaction.

Two HSFs have been identified in human cells, HSF 1 and HSF 2, which bind to the same HSEs and have 38% sequence identity. These factors are activated by distinct stimuli, HSF 1 is responsive to classical stress signals such as heat, heavy metals and oxidative reagents, whereas HSF 2 is activated during hemin-mediated differentiation of human erythroleukemia cells.

HSF 1 exists constitutively in the cytoplasm and the nucleus of unstressed cells as a monomer which lacks DNA binding activity. Through an unknown signal generated during stress, HSF 1 becomes activated to a nuclear localized, trimeric state which binds to DNA. The phosphorylation of HSF 1 is necessary for maximal transcription of heat shock genes.

Synonyms:

HSTF1, HSF-1


Product images:


Figure 1. Western blot analysis of extracts from HUVEC cell treated with TNF- α , using HSF1 (pSer307)Antibody AP08015PU-N (Lane 1 and 2).




Figure 2. Immunofluorescence staining of methanol-fixed HeLa cells using HSF1 (pSer307) antibody AP08015PU.