

Product datasheet for AP08062PU-S

Product datasneet for AP06002PO-3

Product data:

Product Type: Primary Antibodies

c Kit (KIT) Rabbit Polyclonal Antibody

Applications: WE

Recommended Dilution: Western Blot (1/500-1/1000).

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

Immunogen: The antiserum was produced against synthesized non-phosphopeptide derived from human

c-Kit around the phosphorylation site of Tyrosine 721 (N-E-YP-M-D).

Specificity: This antibody detects endogenous levels of total c-Kit protein.

Formulation: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol.

State: Aff - Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Immunoaffinity Chromatography.

Conjugation: Unconjugated

Storage: Store the antibody (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: KIT proto-oncogene receptor tyrosine kinase

Database Link: Entrez Gene 3815 Human

P10721



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

c-Kit is a transmembrane tyrosine kinase encoded by the cKit proto oncogene. c-Kit acts to regulate a variety of biological responses including cell proliferation, apoptosis, chemotaxis and adhesion. Ligand binding to the extracellular domain leads to autophosphorylation on several tyrosine residues within the cytoplasmic domain, and activation. Mutations in c-Kit have been found to be important for tumor growth and progression in a variety of cancers including mast cell diseases, gastrointestinal stromal tumor, acute myeloid leukemia, Ewing sarcoma and lung cancer. Phosphorylation at tyrosine 721 of c-Kit allows binding and activation of PI3 kinase.

Synonyms: SCFR, KIT

Product images:




Figure 1. Western blot analysis of extract from A431 cell treated with EGF 200ng/ml, 30min), using c-Kit Antibody AP08062PU (Lane 1 and 2).