

## Product datasheet for **AP08076PU-N**

### alpha 1 Catenin (CTNNA1) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western Blot: 1/500~1/1000.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	The antiserum was produced against synthesized non-phosphopeptide derived from human Alpha-Catenin around the phosphorylation site of Serine 641 (D-D- <i>Sp</i> -D-F).
Specificity:	This antibody detects endogenous levels of total Alpha-Catenin protein.
Formulation:	PBS (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol. State: Aff - Purified State: Liquid purified Ig fraction.
Concentration:	lot specific
Purification:	Immunoaffinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store the antibody (in aliquots) at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	catenin alpha 1
Database Link:	<a href="#">Entrez Gene 1495 Human P35221</a>



[View online »](#)

**Background:**

The distinct peripheral cytosolic proteins, alpha, beta and gamma-catenin (102, 94 and 86 kDa) found in many tissues bind to the conserved cytoplasmic tail domain of the cell-adhesion cadherins. Catenins link E-cadherin to other integral membrane or cytoplasmic proteins and are modulated by Wnt-1 proto-oncogene. They are good candidates for mediating transduction of cell-cell contact positional signals to the cell interior. Absence of alpha-catenin is found in certain tumor cell lines and reduced levels in certain human carcinomas. Beta-catenin binds directly to the cytoplasmic tail of E-cadherin. It binds to the N-terminus of alpha-catenin and interacts with the protein product of the tumor suppressor gene APC. This interaction involves a 15-aa repeat in the APC. Beta-catenin cell levels seem to be controlled by APC. The central core region of beta-catenin is involved in mediation of cadherin-catenin complex interaction with EGFR.

**Synonyms:**

Alpha E-catenin, Alpha Catenin


**Product images:**

Figure 1. Western blot analysis of extract from A431 cells, untreated or treated with EGF (200ng/ml, 30min), using, Alpha-Catenin Antibody and Alpha-Catenin (Phospho-Ser641) Antibody.