

Product datasheet for AP08252PU-N

AIRE (C-term) Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IHC, R

Recommended Dilution: **ELISA:** 1/32000.

Immunohistochemistry on Paraffin Sections: 2.5 µg/ml.

RIA: 0.01 - 0.03 µg/ml.

Reactivity: Human Host: Goat Isotype: lgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to the C-Terminal of Human AIRE (NP_000374.1,

NP 000649.1).

Epitope: aa534-545

Specificity: This antibody recognize two of the tree reported isoforms (NP_000374.1 and NP_000649.1)

from the C-terminus of human AIRE.

Formulation: Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Immunoaffinity Chromatography

Conjugation: Unconjugated

Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: autoimmune regulator Database Link: Entrez Gene 326 Human

043918

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

AIRE is a transcriptional regulator that is thought to be important in controlling tissue-specific expression of genes in the thymus. The precise function of the protein is not well defined. It contains zinc finger motifs and isoform 1 is localized to both the nucleus and cytoplasm. Three splice variant mRNAs products have been described. The longer AIRE1 mRNA appears to be more abundant and includes exons 1 through 14. Splice variant AIRE2 includes a portion of the non-coding region of exon 1, an alternatively spliced longer exon 8, plus exons 9 through 14. Variant AIRE3 includes the same exon 1-8-9 sequences as found in AIRE2 but utilizes additional alternative splicing in exon 10 that shifts the reading frame such that a stop codon in exon 12 is utilized. The resulting protein products of these splice variants differ significantly. Defects in this gene cause the autosomal-recessive systemic autoimmune disease termed autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED).

Synonyms: Autoimmune regulator

Product images:



Thymus: Formalin-Fixed Paraffin-Embedded (FFPE)