

Product datasheet for AP09067HR-S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Alkaline phosphatase / ALPI / IAP Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IHC, WB

Recommended Dilution: ELISA: 1/10,000-1/50,000.

Western blot: 1/1,000-1/5,000.

Immunohistochemistry.

Reactivity: Bovine
Host: Goat

Isotype: IgG
Clonality: Polyclonal

Immunogen: Alkaline phosphatase from calf intestine

Specificity: This antibody detects bovine alkaline phosphatase. Cross reactivity against

alkaline phosphatase from other tissues and species may occur but have not been specifically

determined.

Immunoelectrophoresis gives a single precipitin arc against anti-Peroxidase, anti-goat serum

as well as purified and partially purified alkaline phosphatase [calf intestine].

Formulation: 0.02M Potassium phosphate, 0.15M Sodium chloride, pH 7.2

Label: HRP

State: Lyophilized purified IgG fraction

Stabilizer: 10 mg/ml BSA (immunoglobulin and protease free)

Preservative: 0.01% (w/v) Gentamicin sulfate (Do NOT add Sodium azide!)

Label: Horseradish peroxidase

Reconstitution Method: Restore with 0.1 ml of deionized water (or equivalent).

Concentration: lot specific

Purification: multi-step process including delipidation, salt fractionation and ion exchange

chromatography followed by extensive dialysis against the buffer

Conjugation: HRP

Storage: Store lyophilized at 2-8°C for 6 months or at -20°C long term.

After reconstitution store the antibody undiluted at 2-8°C for one month or (in alignots) at -

20°C long term. Avoid repeated freezing and thawing.





Alkaline phosphatase / ALPI / IAP Goat Polyclonal Antibody - AP09067HR-S

Stability: Shelf life: one year from despatch.

Database Link: P19111

Background: Alkaline phosphatase (ALP) removes phosphate groups from the 5' end of DNA and RNA, and

from proteins, at high pH. Most mammals have 4 different isozymes: placental, placental like, intestinal and non tissue specific (found in liver, kidney and bone). Tissues with particularly high concentrations of ALP include the liver, bile ducts, placenta, and bone. Damaged or diseased tissue releases enzymes into the blood, so serum ALP measurements can be

abnormal in many conditions, including bone disease and liver disease.

Synonyms: Intestinal alkaline phosphatase