

## Product datasheet for **AP09203AP-N**

### FITC Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ID, IHC, WB
Recommended Dilution:	Suitable for immunoblotting (Western or dot blot), ELISA and immunohistochemistry as well as other phosphatase-antibody based enzymatic assays requiring lot-to-lot consistency. This product is designed for Immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. <u>Recommended Dilutions:</u> ELISA: 1/2,000-1/10,000. Western Blot: 1/500-1/2,500. Immunohistochemistry: 1/200-1/1,000.
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fluorescein conjugated to Goat IgG
Specificity:	This product was prepared from monospecific antiserum by Immunoaffinity chromatography using Fluorescein coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine) and Fluorescein conjugated IgG.
Formulation:	0.05M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 containing 10 mg/ml BSA (IgG and Protease Free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative Label: AP State: Liquid (sterile filtered) purified Ig Label: Alkaline Phosphatase (Calf Intestine) (Molecular Weight 140,000 daltons)
Concentration:	lot specific
Purification:	Affinity Chromatography
Conjugation:	AP



[View online »](#)

<b>Storage:</b>	Store the antibody undiluted at 2-8°C. <b>DO NOT FREEZE!</b>
<b>Stability:</b>	Shelf life: one year from despatch.
<b>Background:</b>	Fluorescein is a fluorophore commonly used to label proteins - protein-fluorescein conjugates are not usually susceptible to precipitation. In addition to its relatively high absorptivity, excellent fluorescence quantum yield and good water solubility, fluorescein has an excitation maximum of 494 nm that closely matches the 488 nm spectral line of the argon-ion laser, making it an important fluorophore for confocal laser-scanning microscopy and flow cytometry applications. Its fluorescence is pH sensitive and is significantly reduced below pH 7. Fluorescein emits most strongly between 500 and 550 nm, but it has a relatively broad emission spectrum reaching to over 600 nm. Several derivatives of fluorescein are commonly used, including FITC (fluorescein isothiocyanate), carboxylates and succinimidyl esters.
<b>Synonyms:</b>	Fluorescein Isothiocyanate