

FITC Goat Polyclonal Antibody

Product datasheet for AP09203TR-N

Product data:

Product Type: Primary Antibodies

Applications: FC

Recommended Dilution: Flow Cytometry.

Immunofluorescence.

Host: Goat Isotype: IgG

Clonality: Polyclonal

Immunogen: Fluorescein conjugated to Goat IgG

Specificity: This antibody reacts to Fluorescein.

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine

Serum Albumin (BSA) as stabilizer and 0.01% (w/v) Sodium Azide as preservative

Label: Texas Red

State: Lyophilized purified Ig

Label: (TM) Sulfonyl Chloride (Molecular Weight 625 daltons)

Absorption emission: 596 nm / 620 nm

Molar radio: 2.9 moles Texas Red(TM) per mole of Goat IgG

Reconstitution Method: Restore with 1,0 ml of deionized water (or equivalent).

Concentration: lot specific

Purification: Affinity chromatography

Conjugation: Texas Red

Storage: Prior to reconstitution store at 2-8°C.

Following reconstitution store the antibody undiluted at 2-8°C for one month

or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



FITC Goat Polyclonal Antibody - AP09203TR-N

Background:

Fluorescein is a fluorophore commonly used to label proteins - protein-fluorescein conjugates are not usually susceptible to precipitation. In addition to its relatively high absorptivity, excellent fluorescence quantum yield and good water solubility, fluorescein has an excitation maximum of 494 nm that closely matches the 488 nm spectral line of the argonion laser, making it an important fluorophore for confocal laser-scanning microscopy and flow cytometry applications. Its fluorescence is pH sensitive and is significantly reduced below pH 7. Fluorescein emits most strongly between 500 and 550 nm, but it has a relatively broad emission spectrum reaching to over 600 nm. Several derivatives of fluorescein are commonly used, including FITC (fluorescein isothiocyanate), carboxylates and succinimidyl esters.

Synonyms:

Fluorescein Isothiocyanate