

Product datasheet for **AP09278PU-N**

CDC27 pSer427 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	ELISA: 1/5,000 - 1/25,000. Western Blot: 1/500 - 1/2,500.
Reactivity:	Canine, Chimpanzee, Chicken, Human, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide corresponding aa 422-430 of Human CDC27
Specificity:	This antibody is directed against the phosphorylated form of CDC27 at the pS427 residue.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 0.01% (w/v) Sodium Azide State: Aff - Purified State: Liquid purified Ig
Concentration:	lot specific
Purification:	Affinity chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	cell division cycle 27
Database Link:	Entrez Gene 996 Human P30260



[View online »](#)

Background:

Human CDC27 (also called Cell division cycle protein 27 homolog, CDC27Hs and H-NUC) shares strong similarity with *Saccharomyces cerevisiae* protein Cdc27. This protein is a component of anaphase-promoting complex (APC), which is composed of eight protein subunits and highly conserved in eukaryotic cells. APC catalyzes the formation of a cyclin B-ubiquitin conjugate that is responsible for the ubiquitin-mediated proteolysis of B-type cyclins. This protein and 3 other members of the APC complex contain the TPR (tetratricopeptide repeat), a protein domain important for protein-protein interaction. This protein was shown to interact with mitotic checkpoint proteins including Mad2, p55CDC and BUBR1, and thus may be involved in controlling the timing of mitosis.

Synonyms:

CDC27Hs, H-NUC, D0S1430E, D17S978E