

Product datasheet for **AP09429RP-N**

Biotin (F(ab')₂ fragment) Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, FC, IF, WB
Recommended Dilution:	Immunofluorescence. Flow Cytometry: 1/100 - 1/250. Western Blot: 1/100-1/250. ELISA: 1/100-1/250.
Host:	Goat
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Biotin conjugated to Keyhole Limpet Hemocyanin (KLH)
Specificity:	This is a Phycoerythrin Conjugated F(ab) ₂ fragment of Anti-Biotin.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA) and 0.01% (w/v) Sodium Azide Label: PE State: Liquid F(ab) ₂ fragment Label: R-Phycoerythrin (Molecular Weight 240,000 daltons) Absorption emission: 488 nm / 575 nm
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity chromatography
Conjugation:	PE
Storage:	Store the antibody undiluted at 2-8°C. DO NOT FREEZE! Store reagent in the dark.
Stability:	Shelf life: 6 month from despatch.



[View online »](#)

Background:

Biotin is a water soluble vitamin, generally classified as a B complex vitamin, also called vitamin B4. After the initial discovery of biotin, nearly forty years of research were required to establish it as a vitamin. Biotin is required by all organisms but can only be synthesized by bacteria, yeasts, molds, algae, and some plant species. Biotin is required as prosthetic group of enzymes involved in incorporation of carbon dioxide into organic compounds. Biotin has a MW of 244 Da.

Synonyms:

Vitamin B7, Vitamin H