

Product datasheet for AP09470PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

c-Myc (MYC) pSer62 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF

Recommended Dilution: Immunofluorescence: 1/100~1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human Myc

around the phosphorylation site of Serine 62(P-L-Sp-P-S).

Specificity: Antibody AP09470PU detects endogenous levels of Myc only when phosphorylated at Serine

62.

Formulation: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol.

State: Aff - Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Immunoaffinity Chromatography using epitope-specific phosphopeptide.

Conjugation: Unconjugated

Storage: Store the antibody (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: v-myc avian myelocytomatosis viral oncogene homolog

Database Link: Entrez Gene 17869 MouseEntrez Gene 24577 RatEntrez Gene 4609 Human

P01106



Background:

The c-Myc protein is a transcription factor, which is encoded by the c-Myc gene on human chromosome 8q24. c-Myc is commonly activated in a variety of tumor cells and plays an important role in cellular proliferation, differentiation, apoptosis and cell cycle progression. The phosphorylation of c-Myc has been investigated and previous studies have suggested a functional association between phosphorylation at Thr58/Ser62 by glycogen synthase kinase 3, cyclin dependent kinase, ERK2 and C-Jun N terminal Kinase (JNK) in cell proliferation and cell cycle regulation. Studies also have shown that c-Myc is essential for tumor cell development in vasculogenesis and angiogenesis that distribute blood throughout the cells, and which brought extensive attention in the development of new therapeutic approach for cancer treatment.

Synonyms:

Transcription factor p64, BHLHE39

Protein Families:

Druggable Genome, Embryonic stem cells, Induced pluripotent stem cells, Stem cell - Pluripotency, Stem cell relevant signaling - JAK/STAT signaling pathway, Stem cell relevant signaling - TGFb/BMP signaling pathway, Stem cell relevant signaling - Wnt Signaling pathway, Transcription Factors

Protein Pathways:

Acute myeloid leukemia, Bladder cancer, Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, ErbB signaling pathway, Jak-STAT signaling pathway, MAPK signaling pathway, Pathways in cancer, Small cell lung cancer, TGF-beta signaling pathway, Thyroid cancer, Wnt signaling pathway

Product images:




Figure 1. Immunofluorescence staining of methanol-fixed HeLa cells using Myc (Phospho-Ser62) Antibody (Red).