

Product datasheet for AP09480PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

TrkA (NTRK1) pSer791 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF

Recommended Dilution: Immunofluorescence: 1/100~1/200.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human TrkA

around the phosphorylation site of tyrosine791 (P-V-Yp-L-D).

Specificity: Antibody AP09480PU detects endogenous levels of TrkA only when phosphorylated at

Tyrosine 791.

Formulation: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol.

State: Aff - Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Immunoaffinity Chromatography using epitope-specific phosphopeptide.

Conjugation: Unconjugated

Storage: Store the antibody (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: neurotrophic receptor tyrosine kinase 1

Database Link: Entrez Gene 4914 Human

P04629

TrkA (NTRK1) pSer791 Rabbit Polyclonal Antibody - AP09480PU-N

Background: The Trk proto-oncogene family contains four members, TrkA, TrkB, TrkC, and TrkE, which are

variably expressed throughout the central and peripheral nervous systems. TrkA binds to nerve growth factor (NGF) and autophosphorylates on tyrosine residues (Tyr490, Tyr674,

Tyr675, Tyr751 and Tyr785) to activate multiple downstream effector proteins.

Phosphorylation at Tyr490 is required for Shc association and subsequent activation of the Ras-MAP kinase-signaling cascade, which leads to activation of Elk-1-dependent gene transcription and neurite growth. Phosphorylations at Tyr674 and Tyr675 lie within the catalytic domain of TrkA tyrosine kinase and reflect Trk kinase activity. Additionally, phosphorylation at Tyr751 is required for PI3-kinase association and activation of the Akt

signaling cascade.

Synonyms: NTRK1, TRK, Trk-A

Protein Families: Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways: Apoptosis, Endocytosis, MAPK signaling pathway, Neurotrophin signaling pathway, Pathways

in cancer, Thyroid cancer

Product images:




Figure 1. Immunofluorescence staining of methanol-fixed HeLa cells using TrkA pSer791 Antibody (Red).