

Product datasheet for AP09684PU-N

Clenbuterol Sheep Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA

Recommended Dilution: ELISA: 10µg/ml.

Host: Sheep lsotype: lgG

Clonality: Polyclonal

Immunogen: Clenbuterol (N)-HSA

Specificity: This antibody reacts to Clenbuterol.

Cross-Reactivity: Clenbuterol 100% Mabuterol 114% Bromobuterol 96% Methyl-Clenbuterol 53%

Mapenterol 49%
Salbutamol 3.8%
Terbutaline 1.6%
Cimaterol 1.0%
Pirbuterol 0.9%
Ractopamine 0.2%
Feneterol 0.2%

Formulation: 20mM Phosphate, 150mM Sodium Chloride, pH 7.2 containing 0.09% Sodium Azide as

preservative
State: Ig Fraction

State: Liquid Ig fraction prepared by Caprylic Acid and Ammonium Sulphate precipitation

procedures

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store the antibody at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Clenbuterol Sheep Polyclonal Antibody - AP09684PU-N

Background:

Clenbuterol belongs to the group of agonists. In livestock production clenbuterol improves the meat/fat ratio in fattened animals or accelerate the growth. Up to now agonists have not been authorized as adjuvants for fattening. In addition to its lipolytic and anabolic effect, clenbuterol has a relaxing effect on non-striated musculature on which is based its therapeutic use as an antiasthmatic and a tocolytic agent. When employed as a fattening adjuvant, as compared with the therapeutic use, clenbuterol is administered in a 5 to 10 times higher dose. Therefore, it is possible that clenbuterol residues may lead to a risk for consumers after illegal administration. Using the clenbuterol monocalantibody, it is possible to detect clenbuterol and other agonists in urine, muscle and liver both rapidly and with accuracy.

Clenbuterol is a long acting beta 2 adrenergic agonist. Like other beta 2 agonists, clenbuterol is believed to act by stimulating production of cyclic AMP through the activation of adenyl cyclase. By definition, Beta 2 agonists have more smooth muscle relaxation activity (bronchial, vascular and uterine smooth muscle) versus its cardiac effects (Beta 1).

Synonyms:

1-(4-amino-3, 5-dichlorophenyl)-2-(tert-butylamino)ethanol, Clenbuterolum, Contraspasmin