

Product datasheet for AP11093PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

EHMT1/GLP (EHMT1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: ELISA: 1/1000.

Western Blot: 1/100-1/500.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide between 956~986 amino acids from the Center region of

Human EUHMTASE1

Specificity: This antibody is specific to Human EUHMTASE1 (Center). **Formulation:** PBS containing 0.09% (W/V) Sodium Azide as preservative

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein G Chromatography, eluted with high and low pH buffers and neutralized

immediately, followed by dialysis against PBS

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: euchromatic histone lysine methyltransferase 1

Database Link: Entrez Gene 79813 Human

Q9H9B1





EHMT1/GLP (EHMT1) Rabbit Polyclonal Antibody - AP11093PU-N


Background:

EHMT1, also known as EUHMTASE1, is a histone methyltransferase. This protein methylates 'Lys-9' of histone H3 in vitro. H3 'Lys-9' methylation represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. EHMT1 is Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. During G0 phase, it probably contributes to silencing of MYC- and E2F-responsive genes, suggesting a role in the G0/G1 transition of the cell cycle.

Synonyms:

EUHMTASE1, GLP, KIAA1876, KMT1D, Eu-HMTase1, G9a-like protein 1, GLP, GLP1, H3-K9-HMTase 5

Product images:



Western blot analysis of EUHMTASE (arrow) using EHMT1 Antibody (Center). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the EUHMTASE gene (Lane 2) (Origene Technologies).