

# **Product datasheet for AP11108PU-N**

#### Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

### CN: techsupport@origene.cn

# **KAT5 (C-term) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1/1,000.

Western blot: 1/100-1/500.

Reactivity: Human
Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the C-terminal region of human HTATIP.

**Specificity:** This antibody is specific to HTATIP (C-term).

**Formulation:** PBS containing 0.09% (W/V) Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

**Concentration:** lot specific

**Purification:** Protein G Chromatography, eluted with high and low pH buffers and neutralized

immediately, followed by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: lysine acetyltransferase 5

Database Link: Entrez Gene 10524 Human

Q92993





#### KAT5 (C-term) Rabbit Polyclonal Antibody - AP11108PU-N


Background: HTATIP belongs to the MYST family of histone acetyl transferases (HATs) and was originally

isolated as an HIV-1 TAT-interactive protein. HATs play important roles in regulating chromatin remodeling, transcription and other nuclear processes by acetylating histone and nonhistone proteins. This protein is a histone acetylase that has a role in DNA repair and

apoptosis and is thought to play an important role in signal transduction.

Synonyms: HTATIP, TIP60, PLIP

# **Product images:**



Western blot analysis of anti-HTATIP Pab in CEM cell line tissue lysate (35ug/lane). HTATIP (arrow) was detected using the purified Pab (1:60 dilution).