

Product datasheet for **AP11206PU-N**

PRDM14 (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	ELISA: 1/1,000. Western blot: 1/100-1/500. Immunohistochemistry: 1/50-1/100.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human PRDM14.
Specificity:	This antibody is specific to PRDM14 (PFM11) (N-term).
Formulation:	PBS containing 0.09% (W/V) Sodium Azide as preservative. State: Purified State: Liquid purified Ig fraction.
Concentration:	lot specific
Purification:	Protein G Chromatography, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	PR domain 14
Database Link:	Entrez Gene 63978 Human Q9GZV8

[View online »](#)

Background:


Similar to acetylation and phosphorylation, histone methylation at the N terminal tail has emerged as an important role in regulating chromatin dynamics and gene activity. Histone methylation occurs on arginine and lysine residues and is catalyzed by two families of proteins, the protein arginine methyltransferase family and the SET domain containing methyltransferase family. Five members have been identified in the arginine methyltransferase family. About 27 are grouped into the SET domain family, and another 17 make up the PR domain family that is related to the SET domain family. PRDM14 is part of a family of PR domain genes that are involved in tumorigenesis. It may function as a transcription factor.

Synonyms:


MGC59730; PFM11

Note:

Predicted MW: 64.062 kDa.

Product images:


Western blot analysis of PRDM14 polyclonal antibody in A549 cell lysate. PRDM14 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.