

Product datasheet for **AP11299PU-N**

BID (BH3 Domain) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	ELISA: 1/1,000. Western blot: 1/100-1/500. Immunohistochemistry: 1/50-1/100.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide encoding the BH3 domain of human Bid.
Specificity:	This antibody is specific to Bid (BH3 Domain).
Formulation:	PBS containing 0.09% (W/V) Sodium Azide as preservative. State: Purified State: Liquid purified Ig fraction.
Concentration:	lot specific
Purification:	Protein G Chromatography, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	BH3 interacting domain death agonist
Database Link:	Entrez Gene 637 Human P55957



[View online »](#)

Background:


Bid is a death agonist that heterodimerizes with either agonist BAX or antagonist BCL2. Bid is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH-terminal part translocates to mitochondria where it triggers cytochrome c release.

Synonyms:


p22 BID

Note:

Calculated MW: 21994 Da

Product images:


LEFT : The anti-Bid BH3 domain Pab is used in Western blot to detect Bid BH3 in HL-60 cell lysate. RIGHT: Western blot analysis of anti-hBid-BH3 Pab in mouse lung tissue lysates (35ug/lane). hBid-BH3 (arrow) was detected using the purified Pab.



LEFT: Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. RIGHT: Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with Bid BH3 Domain Antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.