

Product datasheet for AP11315PU-N

BIK (BH3 Domain) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA: 1/1,000.

Western blot: 1/100-1/500.

Immunohistochemistry: 1/50-1/100.

Reactivity: Human, Mouse

Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected selected from the N-terminal region (BH3 domain) of human Bik.

Specificity: This antibody is specific to BIK (BH3 Domain).

Formulation: PBS containing 0.09% (W/V) Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Protein G Chromatography, eluted with high and low pH buffers and neutralized

immediately, followed by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: BCL2 interacting killer

Database Link: Entrez Gene 638 Human

Q13323



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

BIK (BH3 Domain) Rabbit Polyclonal Antibody - AP11315PU-N

Background: The Bik protein is known to interact with cellular and viral survival-promoting proteins, such


as BCL2 and the Epstein-Barr virus in order to enhance programmed cell death. Because its activity is suppressed in the presence of survival-promoting proteins, this protein is suggested as a likely target for antiapoptotic proteins. This protein shares a critical BH3

domain with other death-promoting proteins, BAX and BAK.


Synonyms: Bcl-2-interacting killer, NBK, BP4, BIP1

Note: Calculated MW: 18016 Da

Product images:



Western blot analysis using anti-Bik BH3 domain Pab to detect Bik BH3 in mouse brain tissue lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.