

Product datasheet for **AP11418PU-N**

Stanniocalcin 2 (STC2) (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	ELISA1: 1,000. Western blotting 1:100 - 1:500. Immunohistochemistry: 1:50 - 1:100.
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	KLH conjugated synthetic peptide selected from the N-terminal region of human STC2
Specificity:	This antibody detects STC2 at N-term.
Formulation:	PBS with 0.09% (W/V) sodium azide State: Purified State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Protein A column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	stanniocalcin 2
Database Link:	Entrez Gene 8614 Human O76061



[View online »](#)

Background:


STC2 is a secreted, homodimeric glycoprotein that is expressed in a wide variety of tissues and may have autocrine or paracrine functions. The encoded protein has 10 of its 15 cysteine residues conserved among stanniocalcin family members and is phosphorylated by casein kinase 2 exclusively on its serine residues. Its C-terminus contains a cluster of histidine residues which may interact with metal ions. The protein may play a role in the regulation of renal and intestinal calcium and phosphate transport, cell metabolism, or cellular calcium/phosphate homeostasis. Constitutive overexpression of human stanniocalcin 2 in mice resulted in pre- and postnatal growth restriction, reduced bone and skeletal muscle growth, and organomegaly. Expression is induced by estrogen and altered in some breast cancers.

Synonyms:


Stanniocalcin-2, STC-2, Stanniocalcin-related protein, STC-related protein, STCRP

Note:

Molecular weight: 33249 Da

Product images:


Western blot analysis of anti-STC2 Antibody (N-term) in mouse kidney tissue lysates (35 ug/lane). STC2 (arrow) was detected using the purified polyclonal antibody.



Formalin-fixed and paraffin-embedded human breast carcinoma tissue reacted with STC2 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.