

Product datasheet for **AP11552PU-N**

DKK2 (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	ELISA 1:1,000. Western blot 1:100 - 1:500. Immunohistochemistry 1:50 - 1:100.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human DKK2.
Specificity:	This antibody detects DKK2 at N-term.
Formulation:	PBS with 0.09% (W/V) sodium azide State: Purified State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS
Conjugation:	Unconjugated
Storage:	Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	dickkopf WNT signaling pathway inhibitor 2
Database Link:	Entrez Gene 27123 Human Q9UBU2



[View online »](#)

Background:


The 259-amino acid DKK2 protein, like DKK1, DKK3, and DKK4, possesses an N-terminal signal peptide and 2 conserved cysteine-rich domains, which are separated by a linker region and contain 10 cys residues each. The second cys region has a putative lipid-binding function that may facilitate WNT/DKK interactions at the plasma membrane. The linker region contains 50 to 55 amino acids in DKK1, DKK2, and DKK4, whereas in DKK3 it contains only 12 amino acids. All DKCs have several potential sites for cleavage by furin-type proteases. Northern blot analysis revealed expression of 4.0- and 4.5-kb DKK2 transcripts in heart, brain, skeletal muscle, and lung. Western blot analysis showed that DKK2 is secreted as a 15- to 17-kD protein. Functional analysis determined that DKK2 does not block *Xenopus* Wnt8 induction of a secondary axis in frog embryos.

Synonyms:


Dickkopf-related protein 2, Dkk-2

Note:


Molecular weight: 28447 Da

Product images:


The anti-DKK2 is used in Western blot to detect DKK2 in Jurkat cell lysate.



Western blot analysis of DKK2 (arrow) using DKK2 Antibody (N-term). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the DKK2 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.