

Product datasheet for AP11586PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

GJA9 (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA 1:1,000.

Immunohistochemistry 1:50 - 1:100.

Reactivity: Human
Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the C-terminal region of human GJA10.

Specificity: This antibody detects GJA10 at C-Term.

Formulation: PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid Ig fraction

Concentration: lot specific

Purification: Protein G column, eluted with high and low pH buffers and neutralized immediately, followed

by dialysis against PBS

Conjugation: Unconjugated

Storage: Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid

repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: gap junction protein alpha 9 **Database Link:** Entrez Gene 81025 Human

P57773

Background: GJA8 is a an integral membrane protein that belongs to the connexin family, alpha-type

(group II) subfamily. One gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from

one cell to a neighboring cell. A connexon is composed of a hexamer of connexins.







Synonyms: Gap junction alpha-9 protein, Connexin-59, Cx59, GJA10, Connexin-58

Note: MOlecular weight: 58856 Da

Product images:



Western blot analysis of GJA10 Antibody (C-term) in mouse heart tissue lysates (35 ug/lane). GJA10 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.