

## Product datasheet for **AP11608PU-N**

### CSEN (KCNIP3) (N-term) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IF, IHC, WB
Recommended Dilution:	<b>ELISA:</b> 1/1,000. <b>Western blot:</b> 1/1,000. <b>Immunofluorescence:</b> 1/10-1/50. <b>Immunohistochemistry on Paraffin Sections:</b> 1/50-1/100.
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1~30 amino acids from the N-terminal region of Human KCHIP3.
Specificity:	This antibody detects KCHIP3 at N-term.
Formulation:	PBS with 0.09% (W/V) Sodium Azide State: Purified State: Liquid purified Ig fraction
Concentration:	lot specific
Purification:	Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	potassium voltage-gated channel interacting protein 3
Database Link:	<a href="#">Entrez Gene 30818 Human Q9Y2W7</a>



[View online »](#)

**Background:**


KCHIP3 is a member of the family of voltage-gated potassium (Kv) channel-interacting proteins (KCNIPs), which belong to the recoverin branch of the EF-hand superfamily. Members of the KCNIP family are small calcium binding proteins. They all have EF-hand-like domains, and differ from each other in the N-terminus. They are integral subunit components of native Kv4 channel complexes. They may regulate A-type currents, and hence neuronal excitability, in response to changes in intracellular calcium. This protein is also shown to function as a calcium-regulated transcriptional repressor, and to interact with presenilins. Mutations in the presenilin genes have been implicated in Alzheimer's disease. Due to utilization of an alternate in-frame translation start codon, the gene for this protein encodes two isoforms with different sizes.

**Synonyms:**


KCNIP3, CSEN, DREAM, KCHIP3, DRE-antagonist modulator

**Note:**


Calculated Molecular weight: 29231 Da

**Product images:**


Western blot analysis of KCHIP3 Antibody (N-term) in mouse heart tissue lysates (35 ug/lane). KCHIP3 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Formalin-fixed and paraffin-embedded human brain tissue reacted with KChIP3 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.