

Product datasheet for AP11708PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

FUBP3 (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA 1:1,000.

Western blot 1:100 - 1:500.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the N-terminal region of human FUBP3.

Specificity: This antibody detects FUBP3 at N-term.

Formulation: PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid Ig fraction

Concentration: lot specific

Purification: Protein G column, eluted with high and low pH buffers and neutralized immediately, followed

by dialysis against PBS

Conjugation: Unconjugated

Storage: Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid

repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: far upstream element binding protein 3

Database Link: Entrez Gene 8939 Human

Q96I24



FUBP3 (N-term) Rabbit Polyclonal Antibody - AP11708PU-N

Background: The far upstream element-binding proteins FUBP, FUBP2, and FUBP3 comprise a family of

single-strand DNA-binding proteins that possess all of the general features of more conventional transcription factors. The FUBPs each bind to a single sequence-specific strand of the far upstream element (FUSE; originally identified upstream of c-myc), and each possesses potent activation domains when fused to the GAL4 DNA-binding domain and assayed by transient transfection. These proteins have also been reported to bind RNA and

participate in various steps of RNA processing, transport or catabolism.

Synonyms: FBP3, FUSE-binding protein 3
Note: Molecular weight: 61640 Da

Product images:



Western blot analysis of anti-FUBP3 in HepG2 cell line lysate (35 ug/lane). FUBP3 (arrow) was detected using the purified Pab (1:60 dilution).