

Product datasheet for AP11835PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Zfp219 (ZNF219) (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA 1:1,000.

Western blot 1:50 - 1:100.

Immunohistochemistry 1:10 - 1:50.

Reactivity: Human
Host: Rabbit
Isotype: Ig

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the N-terminal region of human ZNF219.

Specificity: This antibody detects ZNF219 at N-term.

Formulation: PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid Ig fraction

Concentration: lot specific

Purification: Prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS

Conjugation: Unconjugated

Storage: Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid

repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: zinc finger protein 219

Database Link: Entrez Gene 51222 Human

Q9P2Y4

Background: ZNF219 is a member of the Kruppel-like zinc finger family that is involved in a diverse range

of biological processes. ZNF219 is a 77-kDa nuclear protein containing nine sets of C2H2 zinc

finger structures. It may be involved in transcriptional regulation, it functions as a

transcriptional repressor for the HMGN1 promoter




Zfp219 (ZNF219) (N-term) Rabbit Polyclonal Antibody - AP11835PU-N


Synonyms: Zinc finger protein 219

Note: Molecular weight: 76877 Da

Product images:



Western blot analysis of anti-ZNF219 Antibody (Nterm) in 293 cell line lysates (35 ug/lane). ZNF219 (arrow) was detected using the purified Pab (1:60 dilution).



Formalin-fixed and paraffin-embedded human testis tissue reacted with ZNF219 antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.