

## **Product datasheet for AP11861PU-N**

## Product datasneet for APT 180 IPO-N

## BMPR1B (N-term) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1/1,000.

Western blotting: 1/100-1/500.

Flow Cytometry.

Reactivity: Human, Mouse

**Host:** Rabbit

**Isotype:** lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the N-terminal region of human BMPR1B.

**Specificity:** This antibody recognizes BMPR1B (N-term).

**Formulation:** PBS with 0.09% (W/V) Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

**Concentration:** lot specific

**Purification:** Protein G Chromatography, eluted with high and low pH buffers and neutralized

immediately, followed by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** bone morphogenetic protein receptor type 1B

Database Link: Entrez Gene 658 Human

O00238



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com Background:

The bone morphogenetic protein (BMP) receptors are a family of transmembrane serine/threonine kinases that include the type I receptors BMPR1A and BMPR1B and the type II receptor BMPR2. These receptors are also closely related to the activin receptors, ACVR1 and ACVR2. The ligands of these receptors are members of the TGF-beta superfamily. TGF-betas and activins transduce their signals through the formation of heteromeric complexes with 2 different types of serine (threonine) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II receptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding.

Synonyms: ALK6

Note: Calculated Molecular weight: 56930 Da

## **Product images:**



Western blot analysis of anti-BMPR1B Pab in NCI-H460 cell lysate (Lane A) and mouse heart tissue lysate (Lane B). BMPR1B (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.