

Product datasheet for AP11913PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cripto1 (TDGF1) (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA 1:1,000.

Western blot 1:100 - 1:500.

Immunohistochemistry 1:50 - 1:100.

Reactivity: Human
Host: Rabbit
Isotype: Ig

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the N-terminal region of human TDGF1.

Specificity: This antibody detects CRIPTO (TDGF1) at N-term.

Formulation: PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid Ig fraction

Concentration: lot specific

Purification: Protein G column, eluted with high and low pH buffers and neutralized immediately, followed

by dialysis against PBS

Conjugation: Unconjugated

Storage: Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid

repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: teratocarcinoma-derived growth factor 1

Database Link: Entrez Gene 6997 Human

P13385



Cripto1 (TDGF1) (N-term) Rabbit Polyclonal Antibody - AP11913PU-N


Background: TDGF1 is expressed both in ES cells and during the early phases of embryo development,

while in adults it is reactivated in a wide range of epithelial cancers. This protein could play a role in the determination of the epiblastic cells that subsequently give rise to the mesoderm. It is preferentially expressed in gastric and colorectal carcinomas compared to their normal


counterparts.

Synonyms: Cripto-1 growth factor, CRGF
Note: Molecular weight: 21168 Da

Product images:



The anti-TDGF1 N-term is used in Western blot to detect TDGF1 in A375 cell lysate.



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.