

Product datasheet for AP12039PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

USP21 (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA: 1/1,000.

Western Blot: 1/100-1/500.

Immunohistochemistry: 1/50-1/100.

Reactivity: Human Host: Rabbit

Isotype: Ig

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the C-terminal region of human USP21.

Specificity: This antibody is specific to USP21 (C-term).

Formulation: PBS with 0.09% (W/V) Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Protein G Chromatography, eluted with high and low pH buffers and neutralized

immediately, followed by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: ubiquitin specific peptidase 21

Database Link: Entrez Gene 27005 Human

Q9UK80



USP21 (C-term) Rabbit Polyclonal Antibody - AP12039PU-N

Background: USP21 is a ubiquitin-specific protease, an enzyme that removes ubiquitin from ubiquitinated


proteins. The encoded protein belongs to the C19 peptidase family, also known as family 2 of ubiquitin carboxyl-terminal hydrolases. This protein has been reported to be capable of

removing NEDD8 from NEDD8 conjugates.


Synonyms: USP23

Note: Predicted Molecular weight: 62656 Da

Product images:



Western blot analysis of anti-USP21 Antibody (Cterm) in T47D cell line lysates (35ug/lane). USP21 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.