

Product datasheet for **AP12103PU-N**

STAM1 (STAM) (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	ELISA: 1/1,000. Western Blot: 1/100-1/500. Immunohistochemistry: 1/50-1/100.
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human STAM.
Specificity:	This antibody is specific to STAM (C-term).
Formulation:	PBS with 0.09% (W/V) Sodium Azide as preservative. State: Purified State: Liquid purified Ig fraction.
Concentration:	lot specific
Purification:	Protein G Chromatography, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	signal transducing adaptor molecule
Database Link:	Entrez Gene 8027 Human Q92783



[View online »](#)

Background:


Stimulation of cells with cytokines initiates a signal transduction cascade involving cytokine receptors, Janus kinases (JAKs) and signal transducers and activators of transcription (STATs). STAM for 'signal-transducing adaptor molecule, induced after stimulation of cells with cytokine IL2, is a component of signal transduction downstream of JAK3.1 Human STAM cDNA cloned from a T-cell cDNA library encodes a 540-amino acid protein precipitated by anti-phosphotyrosine. Northern blot analysis indicates that STAM is expressed as a 2.9-kb message in a wide variety of tissue and cell types. The STAM sequence contains a Src-homology 3 (SH3) domain and an immunoreceptor tyrosine-based activation motif (ITAM). It has been suggested that STAM acts as an adaptor molecule in signal transduction pathways from cytokine receptors.

Synonyms:


STAM-1, STAM, Signal transducing adapter molecule 1

Note:

Predicted Molecular weight: 59179 Da

Product images:


(LEFT) The anti-PLIC1 Pab (C-term) is used in Western blot to detect STAM in mouse liver tissue lysate. (RIGHT) Western blot analysis of anti-STAM Antibody (C-term) in HeLa cell line lysates (35ug/lane). STAM (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.