

## Product datasheet for AP12129PU-N

## ARL 1 (ARL1) (N-term) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: ELISA: 1/1,000.

Immunohistochemistry: 1/50-1/100.

Reactivity: Human

Host: Rabbit

Isotype:

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the N-terminal region of human ARL1.

**Specificity:** This antibody is specific to ARL1 (N-term).

Formulation: PBS with 0.09% (W/V) Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Protein G Chromatography, eluted with high and low pH buffers and neutralized immediately,

followed by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2–8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: ADP ribosylation factor like GTPase 1

Database Link: Entrez Gene 400 Human

P40616



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com CN: techsupport@origene.cn




Background:

ARL1 belongs to the ARL (ADP-ribosylation factor-like) family of proteins, which are structurally related to ADP-ribosylation factors (ARFs). ARFs, described as activators of cholera toxin (CT) ADP-ribosyltransferase activity, regulate intracellular vesicular membrane trafficking, and stimulate a phospholipase D (PLD) isoform. Although, ARL proteins were initially thought not to activate CT or PLD, later work showed that they are weak stimulators of PLD and CT in a phospholipid dependent manner.

Synonyms: ARFL1

Note: Predicted Molecular weight: 20417 Da

## **Product images:**



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.