

# Product datasheet for AP12200PU-N

## **FNTA Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

WB **Applications:** 

Recommended Dilution: ELISA: 1/1,000.

Western blot: 1/100-1/500.

Reactivity: Human, Mouse

Host: Rabbit

Isotype: lg

Clonality: Polyclonal

This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide Immunogen:

selected from the center region of human FNTA.

Specificity: This antibody detects FNTA (Center).

Formulation: PBS with 0.09% (W/V) Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

**Purification:** Protein G Chromatography, eluted with high and low pH buffers and neutralized

immediately, followed by dialysis against PBS.

Conjugation: Unconjugated

Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: farnesyltransferase, CAAX box, alpha

Database Link: Entrez Gene 2339 Human

P49354



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

### FNTA Rabbit Polyclonal Antibody - AP12200PU-N


Background:

FNTA, also known as CAAX farnesyltransferase (FTase), attaches a farnesyl group from farnesyl pyrophosphate to cysteine residues at the fourth position from the C terminus of proteins that end in the so-called CAAX box, where C is cysteine, A is usually but not always an aliphatic amino acid, and X is typically methionine or serine. This type of posttranslational modification provides a mechanism for membrane localization of proteins that lack a transmembrane domain. This enzyme has the remarkable property of farnesylating peptides as short as four residues in length that conform to the CAAX consensus sequence. FNTA is also a specific cytoplasmic interactor of the transforming growth factor-beta and activin type I receptors. It is likely to be a key component of the signaling pathway which involves p21ras, an important substrate for farnesyltransferase.

Synonyms: CAAX farnesyltransferase subunit alpha, FTase-alpha, GGTase-I-alpha

Note: Molecular weight: 44408 Da

## **Product images:**



The anti-FNTA Pab is used in Western blot to detect FNTA in mouse brain tissue lysate.