

## Product datasheet for **AP12360PU-N**

### S adenosylhomocysteine hydrolase (AHCY) (C-term) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	ELISA: 1/1,000. Western Blot: 1/50-1/100 Immunohistochemistry: 1/10-1/50.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human AHCY.
Specificity:	This antibody detects S-Adenosylhomocysteine Hydrolase (AdoHcyase/AHCY) (C-term).
Formulation:	PBS with 0.09% (W/V) Sodium Azide as preservative. State: Purified State: Liquid purified Ig fraction.
Concentration:	lot specific
Purification:	Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	adenosylhomocysteinase
Database Link:	<a href="#">Entrez Gene 191 Human P23526</a>



[View online »](#)

**Background:**


S-adenosylhomocysteine hydrolase (AHCY) catalyzes the reversible hydrolysis of S-adenosylhomocysteine (AdoHcy) to adenosine (Ado) and L-homocysteine (Hcy). Thus, it regulates the intracellular S-adenosylhomocysteine (SAH) concentration thought to be important for transmethylation reactions. Deficiency in this protein is one of the different causes of hypermethioninemia. S-adenosylhomocysteine hydrolase belongs to the adenosylhomocysteinase family.

**Synonyms:**


Adenosylhomocysteinase, SAHH

**Note:**

**Molecular weight:** 47716 Da

**Product images:**

Western blot analysis of anti-AHCY Antibody (C-term) in HepG2 cell line lysates (35ug/lane). AHCY (arrow) was detected using the purified Pab (1:60 dilution).



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with AHCY antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.