

Product datasheet for AP12458PU-N

Ataxin 1 (ATXN1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1/1,000.

Western Blot: 1/50-1/100.

Reactivity: Human
Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This ATXN1 antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 754-781 amino acids from human ATXN1.

Specificity: This antibody detects Ataxin-1 (ATXN1).

Formulation: PBS with 0.09% (W/V) Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Protein A Chromatography followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: ataxin 1

Database Link: Entrez Gene 6310 Human

P54253



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The autosomal dominant cerebellar ataxias (ADCA) are a heterogeneous group of neurodegenerative disorders characterized by progressive degeneration of the cerebellum, brain stem and spinal cord. Clinically, ADCA has been divided into three groups: ADCA types I-III. ADCAI is genetically heterogeneous, with five genetic loci, designated spinocerebellar ataxia (SCA) 1, 2, 3, 4 and 6, being assigned to five different chromosomes. ADCAII, which always presents with retinal degeneration (SCA7), and ADCAIII often referred to as the `pure' cerebellar syndrome (SCA5), are most likely homogeneous disorders. Several SCA genes have been cloned and shown to contain CAG repeats in their coding regions. ADCA is caused by the expansion of the CAG repeats, producing an elongated polyglutamine tract in the corresponding protein. The expanded repeats are variable in size and unstable, usually increasing in size when transmitted to successive generations. The function of the ataxins is not known.

Synonyms: Ataxin 1, ATXN1, ATX1, SCA1

Note: Molecular weight: 86923 Da

Product images:



Western blot analysis of anti-ATXN1 Antibody (S776) in CEM cell line lysates (35ug/lane). ATXN1 (arrow) was detected using the purified Pab.