

## Product datasheet for AP12546PU-N

## Histone H3 pSer10 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

**Recommended Dilution:** ELISA: 1/1,000.

Western Blot: 1/100-1/500.

Immunohistochemistry: 1/50-1/100.

Reactivity: Human

Host: Rabbit

Isotype:

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic

phosphopeptide corresponding to amino acid residues surrounding S10 of human H3.

**Specificity:** This antibody detects H3 pSer10.

Predicted to cross react with Mouse, Rabbit and Zebrafish (100% Antigen Homology).

Formulation: PBS with 0.09% (W/V) Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Protein G Affinity Chromatography. Then, the antibody fraction is peptide affinity purified in a

2-step procedure with control and phosphorylated peptides. The phospho-specific antibody is eluted with high and low pH buffers and neutralized immediately, followed by dialysis against

PBS.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com CN: techsupport@origene.cn




Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.


**Synonyms:** H3 Histone family, HIST1H3

Note: Molecular weight: 15404 Da

## **Product images:**



Western blot analysis of anti-Phospho-H3 pSerl0 Pab (Cat. #APl2546PU-N) in HepG2 cell line lysate (35ug/lane).Phospho-H3-pSl0 (arrow) was detected using the purified Pab (1:64 dilution).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clini