

## Product datasheet for **AP12900PU-N**

### **E2F1 pSer332 Rabbit Polyclonal Antibody**

#### **Product data:**

|                       |   |
|-----------------------|---|
| Product Type:         | Primary Antibodies  |
| Recommended Dilution: | ELISA: 1/1,000.<br>Dot Blot: 1/500.   |
| Reactivity:           | Human   |
| Host:                 | Rabbit  |
| Isotype:              | Ig  |
| Clonality:            | Polyclonal  |
| Immunogen:            | This antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S332 of human E2F1. |
| Specificity:          | This antibody detects E2F1 pSer332.   |
| Formulation:          | PBS with 0.09% (W/V) Sodium Azide as preservative.<br>State: Aff - Purified<br>State: Liquid purified Ig fraction.  |
| Concentration:        | lot specific  |
| Purification:         | Protein A Chromatography followed by two-step phosphospecific peptide affinity purification.  |
| Conjugation:          | Unconjugated  |
| Storage:              | Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.<br>Avoid repeated freezing and thawing.                                     |
| Stability:            | Shelf life: one year from despatch.   |
| Gene Name:            | E2F transcription factor 1  |
| Database Link:        | <a href="#">Entrez Gene 1869 Human Q01094</a>   |

[View online »](#)

**Background:**


E2F1 is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of the family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can mediate both cell proliferation and p53-dependent/independent apoptosis.

**Synonyms:**

E2F-1, RBBP3, PBR3

**Note:**

**Molecular weight:** 46920 Da

**Product images:**


Dot blot analysis of anti-E2F1-pSer332 Pab (Cat#AP12900PU-N) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

**Dot Blot**