

Product datasheet for AP12997PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

ACK1 (TNK2) pTyr518 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Recommended Dilution: ELISA: 1/1,000.

Dot Blot: 1/500.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic

phosphopeptide corresponding to amino acid residues surrounding Y518 of human ACK1.

Specificity: This antibody detects TNK2 / ACK1 pTyr518.

Formulation: PBS with 0.09% (W/V) Sodium Azide as preservative.

State: Aff - Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Protein A Chromatography followed by two-step phosphospecific peptide affinity purification.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: tyrosine kinase non receptor 2

Database Link: Entrez Gene 10188 Human

Q07912

Background: ACK1 is a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the

intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active

form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal

transduction pathway.




ACK1 (TNK2) pTyr518 Rabbit Polyclonal Antibody - AP12997PU-N

Synonyms: ACK; ACK-1; ACK1; FLJ44758; FLJ45547; p21cdc42Hs

Note: Molecular weight: 114327 Da

Product images:



Dot Blot

Dot blot analysis of anti-Phospho-ACK1 pTyr518 Antibody (Cat.#AP12997PU-N) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per