

Product datasheet for AP13204PU-N

OASL (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

IHC, WB **Applications:**

Recommended Dilution: ELISA: 1/1,000.

Western blotting: 1/100 - 1/500.

Immunohistochemistry: 1/50 - 1/100.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the C-terminal region of human OASL.

Specificity: This antibody reacts to OASL.

Formulation: PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid purified Ig

Concentration: lot specific

Purification: Protein G column, eluted with high and low pH buffers and neutralized immediately, followed

by dialysis against PBS

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: 2'-5'-oligoadenylate synthetase like

Database Link: Entrez Gene 8638 Human

Q15646



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

2-prime,5-prime oligoadenylates (2-5As) bind to and activate RNase L, leading to degradation of RNA and inhibition of protein synthesis. 2-5As are produced by 2-5A synthetases (OASs), a highly-conserved family of interferon-induced enzymes. The predicted 514-amino acid human p59OASL (2-5A synthetases-like) protein shares a highly conserved N-terminal domain with other OASs. The C-terminal portion of p59OASL contains 2 ubiquitin-like domains. p59OASL is expressed in most tissues, with the highest levels in hematopoietic tissues, colon, and stomach.


Synonyms:

p59 OASL, TR-interacting protein 14

Product images:



The anti-OASL C-term Antibody is used in Western blot to detect OASL in HL60 lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining.