

Product datasheet for AP13594PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Pyruvate Kinase (PKLR) (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA: 1/1,000.

Western blotting: 1/100 - 1/500.

Immunohistochemistry: 1/50 - 1/100.

Reactivity: Human, Mouse

Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the C-terminal region of human PKLR.

Specificity: This antibody reacts to PKLR.

Formulation: PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid purified Ig

Concentration: lot specific

Purification: Protein G column, eluted with high and low pH buffers and neutralized immediately, followed

by dialysis against PBS

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: pyruvate kinase, liver and RBC

Database Link: Entrez Gene 5313 Human

P30613





Pyruvate Kinase (PKLR) (C-term) Rabbit Polyclonal Antibody – AP13594PU-N


Background:

PKLR is a pyruvate kinase that catalyzes the production of phosphoenolpyruvate from pyruvate and ATP. Defects in this enzyme, due to gene mutations or genetic variations, are the common cause of chronic hereditary nonspherocytic hemolytic anemia (CNSHA or HNSHA).


Synonyms:

PK1, PKL, Pyruvate kinase R/L, Pyruvate kinase 1, R-type/L-type pyruvate kinase, Red cell/liver pyruvate kinase

Product images:



The anti-PKLR Pab is used in Western blot to detect PKLR in mouse liver tissue lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining.