

Product datasheet for **AP13679PU-N**

ADK (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	ELISA: 1/1,000. Western blotting: 1/100 - 1/500. Flow cytometry.
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human ADK.
Specificity:	This antibody reacts to ADK.
Formulation:	PBS with 0.09% (W/V) sodium azide State: Purified State: Liquid purified Ig
Concentration:	lot specific
Purification:	Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	adenosine kinase
Database Link:	Entrez Gene 132 Human P55263


[View online »](#)

Background:

Adenosine kinase (ATP:adenosine 5-prime-phosphotransferase) is an abundant enzyme in mammalian tissues that catalyzes the transfer of the gamma-phosphate from ATP to adenosine, thereby serving as a potentially important regulator of concentrations of both extracellular adenosine and intracellular adenine nucleotides. Adenosine has widespread effects on the cardiovascular, nervous, respiratory, and immune systems and inhibitors of ADK could play an important pharmacological role in increasing intravascular adenosine concentrations and acting as antiinflammatory agents. The encoded protein does not present any sequence similarities to other well-characterized mammalian nucleoside kinases. In contrast, 2 regions were identified with significant sequence identity to microbial ribokinase and fructokinases and a bacterial inosine/guanosine kinase. Thus, ADK is a structurally distinct mammalian nucleoside kinase that appears to be akin to sugar kinases of microbial origin. Animal studies have demonstrated that a deficiency of adenosine metabolism a powerful contributor to the development of neonatal hepatic steatosis, providing a model for the rapid development of postnatally lethal fatty liver.

Synonyms:

ADK, AK

Product images:


The anti-ADK Pab is used in Western blot to detect ADK in mouse bladder tissue lysate