

Product datasheet for AP13760PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

GUCY1A2 (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA: 1/1,000.

Immunohistochemistry: 1/50 - 1/100.

Western Blot: 1/50 - 1/200.

Reactivity: Human, Mouse

Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the N-terminal region of human GUCY1A2.

Specificity: This antibody reacts to Guanylyl Cyclase alpha 2 (GUCY1A2).

Formulation: PBS with 0.09% (W/V) Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: guanylate cyclase 1, soluble, alpha 2

Database Link: Entrez Gene 2977 Human

P33402

Background: Soluble guanylate cyclase (sGC), a heterodimeric protein consisting of an alpha and a beta

subunit, catalyzes the conversion of GTP to the second messenger cGMP and functions as the

main receptor for nitric oxide and nitrovasodilator drugs.


Synonyms: Guanylate cyclase soluble subunit alpha-2, GUC1A2, GUCSA2






Note: Calculated MW: 81750 Da

Product images:



Western blot analysis of anti-GUCY1A2 Antibody (N-term) in mouse cerebellum tissue lysates (35ug/lane). GUCY1A2 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining.