

Product datasheet for AP13860PU-N

Phone: +1-888-267-4436 https://www.origene.com

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Rockville, MD 20850, US

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

UCK2 (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

IHC, WB **Applications:**

Recommended Dilution: ELISA 1:1,000.

Western Blot 1:100 - 1:250.

Immunohistochemistry 1:50 - 1:100.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the N-terminal region of human UCK2.

Specificity: This antibody detects UCK2 at N-term. Formulation: PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid Ig fraction

Concentration: lot specific

Purification: Protein G column, eluted with high and low pH buffers and neutralized immediately, followed

by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid

repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: uridine-cytidine kinase 2

Database Link: Entrez Gene 7371 Human

Q9BZX2



UCK2 (N-term) Rabbit Polyclonal Antibody - AP13860PU-N

Background: UCK2 catalyzes the phosphorylation of uridine monophosphate to uridine diphosphate. This

is the first step in the production of the pyrimidine nucleoside triphosphates required for RNA


and DNA synthesis. In addition, an allele of this gene may play a role in mediating

nonhumoral immunity to Hemophilus influenzae type B.


Synonyms: UCK 2, Uridine-cytidine kinase 2, TSA903, UMPK

Note: Molecular weight: 29299 Da.

Product images:



Western blot analysis of anti-UCK2 Antibody (N-term) pre-incubated with and without blocking peptide in Hela cell line lysate. UCK2 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue (hepatocarcinoma) reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.