

# Product datasheet for AP13898PU-N

## DCAMKL1 (DCLK1) (C-term) Rabbit Polyclonal Antibody

### **Product data:**

#### **Product Type: Primary Antibodies** IHC, WB **Applications: Recommended Dilution:** ELISA: 1/1.000. Western blot: 1/100-1/500. Immunohistochemistry: 1/10-1/50. **Reactivity:** Human, Mouse Host: Rabbit Isotype: lg **Clonality:** Polyclonal Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human DCAMKL1. This antibody detects DCAMKL1 at C-term. Specificity: Formulation: PBS with 0.09% (W/V) Sodium Azide as preservative. State: Purified State: Liquid purified lg fraction. **Concentration:** lot specific **Purification:** Protein G Chromatography eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS. **Conjugation:** Unconjugated Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage: Avoid repeated freezing and thawing. Stability: Shelf life: one year from despatch. Gene Name: doublecortin like kinase 1 Database Link: Entrez Gene 9201 Human 015075



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	DCAMKL1 (DCLK1) (C-term) Rabbit Polyclonal Antibody – AP13898PU-N
Background:	Doublecortin-like kinase (DCAMKL1)(Ser/Thr protein kinase family) is essential for proper neurogenesis, neuronal migration, and axonal wiring. DCAMKL1 is involved in a calcium- signaling pathway controling neuronal migration in the developing brain, and participates in functions of the mature nervous system. DCAMKL1 protein shares high homology with doublecortin (DCX). DCLK, but not DCX, is highly expressed in regions of active neurogenesis in the neocortex and cerebellum. DCAMKL1 controls mitotic division by regulating spindle formation and also determines the fate of neural progenitors during cortical neurogenesis.
Synonyms:	DCLK, DCKL1, DCAMKL1, DCDC3A, CLICK1, KIAA0369
Note:	Predicted Molecular weight: 82223 Da

## **Product images:**



The DCAMKL1 Antibody (C-term ) is used in Western blot to detect DCAMKL1 in mouse heart tissue lysate.

Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with DCAMKL1 Antibody (C-term), which was peroxidaseconjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US