

## Product datasheet for **AP13936PU-N**

### CDKL5 (C-term) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	ELISA 1:1,000. Western blot 1:100 - 1:500. Immunohistochemistry 1:10 - 1:50.
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human STK9.
Specificity:	This antibody detects CDKL5 (STK9) at C-term.
Formulation:	PBS with 0.09% (W/V) sodium azide State: Purified State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	cyclin dependent kinase like 5
Database Link:	<a href="#">Entrez Gene 6792 Human O76039</a>



[View online »](#)

**Background:**


Defects in STK9, a dual-specificity serine/threonine kinase, are a cause of atypical Rett syndrome. Rett syndrome is an X-linked dominant disease. It is a progressive neurologic developmental disorder and one of the most common causes of mental retardation in females. Patients appear to develop normally until 6 to 18 months of age, then gradually lose speech and purposeful hand movements and develop microcephaly, seizures, autism, ataxia, intermittent hyperventilation, and stereotypic hand movements. After initial regression, the condition stabilizes and patients usually survive into adulthood. Rett syndrome due to CDKL5-associated mutations is characterized by a severe early-onset phenotype and atypical features such as infantile spasms.

**Synonyms:**


ISSX; STK9

**Note:**

Molecular weight: 115537 Da

**Product images:**


The anti-STK9 Pab is used in Western blot to detect STK9 in mouse lung tissue lysate.



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with STK9 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.