

# Product datasheet for AP14369PU-N

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Integrin Linked Kinase (ILK) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: ELISA: 1/1,000.

Immunohistochemistry: 1/10 - 1/50.

Reactivity: Human
Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide corresponding to amino acid residues surrounding Y343 of

human ILK.

**Specificity:** This antibody reacts to ILK.

Formulation: PBS with 0.09% (W/V) sodium azide

State: Aff - Purified State: Liquid purified Ig

**Concentration:** lot specific

**Purification:** Affinity chromatography on Protein A

**Conjugation:** Unconjugated

**Storage:** Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** integrin linked kinase

Database Link: Entrez Gene 3611 Human

Q13418



#### Integrin Linked Kinase (ILK) Rabbit Polyclonal Antibody - AP14369PU-N


Background:

Transduction of extracellular matrix signals through integrins influences intracellular and extracellular functions, and appears to require interaction of integrin cytoplasmic domains with cellular proteins. Integrin-linked kinase (ILK), interacts with the cytoplasmic domain of beta-1 integrin. ILK is a serine/threonine protein kinase with 4 ankyrin-like repeats, which associates with the cytoplasmic domain of beta integrins and acts as a proximal receptor kinase regulating integrin-mediated signal transduction.

Synonyms:

ILK-1, ILK-2, p59ILK, ILK, ILK1, ILK2

# **Product images:**



Formalin-fixed and paraffin-embedded human skeletal muscle tissue reacted with Phospho ILK Antibody (S343), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.