

## Product datasheet for **AP14441PU-N**

### DDR2 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	ELISA: 1/1,000. Western blotting: 1/50 - 1/100.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a his tag recombinant protein of human TYRO10.
Specificity:	This antibody reacts to TYRO10.
Formulation:	PBS with 0.09% (W/V) sodium azide State: Purified State: Purified polyclonal antibody supplied in . This antibody is
Concentration:	lot specific
Purification:	Prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	discoidin domain receptor tyrosine kinase 2
Database Link:	<a href="#">Entrez Gene 4921 Human Q16832</a>




[View online »](#)

**Background:**

Receptor tyrosine kinases (RTKs) play a key role in the communication of cells with their microenvironment. These molecules are involved in the regulation of cell growth, differentiation and metabolism. In several cases the biochemical mechanism by which RTKs transduce signals across the membrane has been shown to be ligand induced receptor oligomerization and subsequent intracellular phosphorylation. This autophosphorylation leads to phosphorylation of cytosolic targets as well as association with other molecules, which are involved in pleiotropic effects of signal transduction. RTKs have a tripartite structure with extracellular, transmembrane and cytoplasmic regions. There are several subclasses of RTKs and TYRO10 belongs to a novel subclass. The deduced amino acid sequence of TYRO10 has a unique extracellular region encompassing a factor VIII-like domain, not previously described for RTKs.

**Synonyms:**

NTRKR3, TKT, TYRO10

**Product images:**

(LEFT) Western blot analysis of anti-TYRO10 Antibody in Jurkat cell line lysates (35ug/lane). TYRO10 (arrow) was detected using the purified Pab. (RIGHT) Western blot analysis of TYRO10 (arrow) using rabbit polyclonal TYRO10 Antibody. 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the DDR2 gene (Lane 2)