

Product datasheet for AP14531PU-N

RND3 (Center) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: ELISA: 1/1,000.

Immunohistochemistry: 1/10 - 1/50.

Reactivity: Human Rabbit Host:

Isotype: lg

Clonality: Polyclonal

This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide Immunogen:

selected from the Center region of human AEHE.

Specificity: This antibody reacts to ARHE.

Formulation: PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid purified Ig

Concentration: lot specific

Purification: Prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS

Conjugation: Unconjugated

Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: Rho family GTPase 3

Database Link: Entrez Gene 390 Human

P61587



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Members of the Rho family of Ras-related GTPases, such as ARHE, regulate the organization of the actin cytoskeleton in response to extracellular growth factors. Like Ras (MIM 190020), Rho family members appear to cycle between an inactive GDP-bound form and an active GTP-bound form. Three major regulators of Rho activity have been identified: RhoGDIs, which interact with the GDP-bound Rho proteins to keep them in a resting complex (see MIM 601925); GEFs, which promote GDP/GTP exchange leading to activation of Rho proteins (see MIM 601855); and GAPs, which stimulate GTP hydrolysis and return the activated Rho protein to its inactive form (see MIM 602680) (Nobes et al., 1998 [PubMed 9531558]).

Synonyms:

RHO8, RHOE, MemB protein

Product images:



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with ARHE antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.