

## **Product datasheet for AP14602PU-N**

## IRAK2 (C-term) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: ELISA: 1/1,000.

Western blotting: 1/100 - 1/500.

Immunohistochemistry: 1/50 - 1/100.

Reactivity: Human
Host: Rabbit

**Isotype:** lg

Clonality: Polyclonal

**Immunogen:** This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the C-terminal region of human IRAK2.

**Specificity:** This antibody reacts to IRAK2.

Formulation: PBS with 0.09% (W/V) sodium azide

State: Purified

State: Liquid purified Ig

**Concentration:** lot specific

**Purification:** Protein G column, eluted with high and low pH buffers and neutralized immediately, followed

by dialysis against PBS

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** interleukin 1 receptor associated kinase 2

Database Link: Entrez Gene 3656 Human

O43187



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:


Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The tyrosine-like kinase (TLK) group consists of 40 tyrosine and serine-threonine kinases such as MLK (mixed-lineage kinase), LISK (LIMK/TESK), IRAK (interleukin-1 receptor-associated kinase), Raf, RIPK (receptor-interacting protein kinase), and STRK (activin and TGF-beta receptors) families.

Synonyms: IRAK-2

## **Product images:**



Western blot analysis of anti-IRAK2 Pab in K562 cell lysate. IRAK2 (Arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.