

Product datasheet for **AP14626PU-N**

LIM Kinase 1 (LIMK1) (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	ELISA: 1/1,000. Western blotting: 1/100 - 1/500. Immunohistochemistry: 1/50 - 1/100.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human LIMK1.
Specificity:	This antibody reacts to LIM Kinase 1 (LIMK1).
Formulation:	PBS with 0.09% (W/V) sodium azide State: Purified State: Liquid purified Ig
Concentration:	lot specific
Purification:	Protein G column, eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	LIM domain kinase 1
Database Link:	Entrez Gene 3984 Human P53667




[View online »](#)

Background:


LIMK1, a member of the Ser/Thr protein kinase family, may be a component of an intracellular signaling pathway and may be involved in brain development. It phosphorylates and inactivates the actin binding/depolymerizing factor cofilin and induces actin cytoskeletal changes. The LIM domain interacts with the cytoplasmic domain of NRG1, and this cytoplasmic protein also binds ROCK1, which phosphorylates LIMK1 on serine and/or threonine residues. Highest expression occurs in both adult and fetal nervous systems. It is detected ubiquitously throughout the different regions of adult brain, with highest levels in the cerebral cortex, and is expressed to a lesser extent in heart and skeletal muscle. Haploinsufficiency of LIMK1 may be the cause of certain cardiovascular and musculo-skeletal abnormalities observed in Williams-Beuren syndrome (WBS), a rare developmental disorder. It is a contiguous gene deletion syndrome involving genes from chromosome band 7q11.23. This protein contains 2 LIM zinc-binding domains and 1 PDZ/DHR domain.

Synonyms:

LIMK, LIM domain kinase 1, LIMK-1, EC=2.7.11.1

Product images:

The anti-LIMK1 Pab is used in Western blot to detect LIMK1 in A375 cell lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.