

Product datasheet for AP14722PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cytochrome P450 2C9 (CYP2C9) (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1/1,000.

Western blot: 1/50-1/100.

Reactivity: Human
Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: This antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide

selected from the N-terminal region of human CYP2C9.

Specificity: This antibody is specific to CYP2C9 (N-term).

Formulation: PBS with 0.09% (W/V) Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: cytochrome P450 family 2 subfamily C member 9

Database Link: Entrez Gene 1559 Human

P11712

Background: CYP2C9 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450

proteins are monooxygenases which catalyze many reactions involved in drug metabolism

and synthesis of cholesterol, steroids and other lipids. This protein localizes to the

endoplasmic reticulum and its expression is induced by rifampin. The enzyme is known to metabolize many xenobiotics, including phenytoin, tolbutamide, ibuprofen and S-warfarin.






Cytochrome P450 2C9 (CYP2C9) (N-term) Rabbit Polyclonal Antibody - AP14722PU-N

Synonyms: Cytochrome P450 2C9, CYP2C10, CYPIIC9, Cytochrome P450 PB-1, Cytochrome P450 MP-

4/MP-8

Note: Calculated MW: 55628 Da

Product images:



Western blot analysis of anti-CYP2C9 Antibody (N-term) in Jurkat cell line lysates (35ug/lane). CYP2C9 (arrow) was detected using the purified Pab.